

**WO9917796**

Publication Title:

**MODULATION OF LERK-2-MEDIATED CELL ADHESION**

Abstract:

Methods of modulating LERK-2-mediated cell adhesion, as well as methods of modulating angiogenesis and inflammation are described. Also described are agents such as antibodies which can modulate LERK-2 mediated cell adhesion, as well as methods of treating angiogenic diseases and inflammatory diseases.

-----  
Data supplied from the esp@cenet database - <http://ep.espacenet.com>

*This Patent PDF Generated by Patent Fetcher(TM), a service of Stroke of Color, Inc.*



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 38/17, 39/395, G01N 33/53, C07K 16/28, C12N 5/20</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 99/17796</b> <b>(43) International Publication Date:</b> 15 April 1999 (15.04.99)
<b>(21) International Application Number:</b> PCT/US98/20557 <b>(22) International Filing Date:</b> 30 September 1998 (30.09.98) <b>(30) Priority Data:</b> 60/060,757 2 October 1997 (02.10.97) US <b>(71) Applicant (for all designated States except US):</b> LEUKOSITE, INC. [US/US]; 215 First Street, Cambridge, MA 02142 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> BRISKIN, Michael, J. [US/US]; 28 Harbell Street, Lexington, MA 02421 (US). ZOU, Lily [CN/US]; 266A Harvard Street No. 5, Cambridge, MA 02139 (US). <b>(74) Agents:</b> WENDLER, Helen, E. et al.; Hamilton, Brook, Smith & Reynolds, P.C., Two Militia Drive, Lexington, MA 02421 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i> <i>With an indication in relation to deposited biological material furnished under Rule 13bis separately from the description.</i>
<b>(54) Title:</b> MODULATION OF LERK-2-MEDIATED CELL ADHESION <b>(57) Abstract</b> <p>Methods of modulating LERK-2-mediated cell adhesion, as well as methods of modulating angiogenesis and inflammation are described. Also described are agents such as antibodies which can modulate LERK-2 mediated cell adhesion, as well as methods of treating angiogenic diseases and inflammatory diseases.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

-1-

## MODULATION OF LERK-2-MEDIATED CELL ADHESION

## RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Application No. 60/060,757, filed October 2, 1997, the entire teachings of which are incorporated herein by reference.

## BACKGROUND OF THE INVENTION

In order to coordinate the activities necessary for survival of a multicellular organism, the individual cells of the organism must communicate with each other via cell-surface receptors and their ligands. The known receptors are generally classified based on the particular method of signal transduction (translation of an extracellular signal into an intracellular signal) which is most characteristic of the receptor class. One class of receptors is the group of receptor protein-tyrosine kinases, which have intracellular kinase domains that are activated by binding of one or more ligands, resulting in autophosphorylation of specific tyrosine residues and activation of intracellular signalling molecules.

The largest subfamily of the receptor protein-tyrosine kinases (RPTK) is the Eph subfamily, characterized by the oncogene *eph* (Hirai et al., *Science* 238:1717-1720 (1987)). The ligands for the Eph RPTKs constitute a rapidly expanding group of molecules which are either transmembrane or glycosylphosphatidylinositol (GPI)-linked (Pandey et al., *Current Biology* 5(9):986-989 (1995)). It is likely that the Eph receptor-ligand

-2-

interactions play important roles in the growth, development and survival of multicellular organisms.

#### SUMMARY OF THE INVENTION

The invention pertains to a method of modulating LERK-2-mediated cell adhesion, comprising the step of contacting a cell expressing LERK-2 and/or a cell expressing a receptor for LERK-2 with an inhibitor or promoter of LERK-2-mediated cell adhesion. In a particular embodiment, the receptor for LERK-2 is selected from the group consisting of Nuk, Cek5, Tyro5 and ERK, and in a preferred embodiment the receptor for LERK-2 is Nuk. In one embodiment, the cell expressing a receptor for LERK-2 is a lymphoid cell, particularly a T cell.

In one embodiment of the method, the inhibitor is selected from the group consisting of an antibody or functional antibody fragment which inhibits LERK-2-mediated cell adhesion (e.g., an anti-LERK-2 antibody), soluble LERK-2 and a LERK-2-Ig chimera. In a particular embodiment, the anti-LERK-2 antibody or fragment is selected from the group consisting of monoclonal antibody 2A1, a functional antibody fragment of 2A1, monoclonal antibody 4A1 and a functional antibody fragment of 4A1.

The invention also relates to a method of detecting or identifying an inhibitor or promoter of LERK-2-mediated cell adhesion, comprising the steps of combining an agent to be tested with a composition comprising a cell expressing LERK-2 and a composition comprising a cell expressing a receptor for LERK-2 under conditions suitable for binding of LERK-2 to a receptor for LERK-2, and detecting or measuring cell adhesion between the cell expressing LERK-2 and the cell expressing a receptor for LERK-2, whereby inhibition or promotion of cell adhesion

-3-

by the agent is indicative that the agent is an inhibitor or promoter, respectively. The invention also pertains to inhibitors and promoters (i.e., modulators) identified by the described methods.

5           For example, modulators which can be identified as described herein include modulating antibodies, e.g., antibodies which inhibit or promote LERK-2-mediated cell adhesion. Modulating antibodies can be an antibody or functional antibody fragment which modulates LERK-2-  
10 mediated cell adhesion, such as an antibody or fragment which modulates binding of LERK-2 to a receptor for LERK-2. In a particular embodiment, the antibody or functional antibody fragment is an anti-LERK-2 antibody.

          In one embodiment the invention relates to an  
15 antibody or functional antibody fragment which inhibits LERK-2-mediated cell adhesion, such as an antibody or fragment which inhibits binding of LERK-2 to a receptor for LERK-2. In a particular embodiment, the antibody or functional antibody fragment is an anti-LERK-2 antibody.  
20 In a preferred embodiment, the antibody or functional antibody fragment is selected from the group consisting of monoclonal antibody 2A1, monoclonal antibody 4A1, a functional antibody fragment of 2A1 and a functional antibody fragment of 4A1. The invention also relates to  
25 an antibody or functional antibody fragment which can compete with monoclonal antibody 2A1 or monoclonal antibody 4A1 for binding to LERK-2 or portion thereof.

          The invention also pertains to the hybridoma cell line deposited under ATCC Accession No. HB-12412 and to  
30 the hybridoma cell line deposited under ATCC Accession No. HB-12413. The invention further relates to a monoclonal antibody or antigen binding fragment thereof produced by

-4-

the hybridoma cell line deposited under ATCC Accession No. HB-12412 or HB-12413.

The invention also relates to a method for treating an angiogenic disease, comprising administering to a mammal a therapeutically effective amount of an inhibitor of LERK-2-mediated cell adhesion, whereby angiogenesis is reduced.

The invention further pertains to a method for treating an inflammatory disease or condition, comprising administering to a mammal a therapeutically effective amount of an inhibitor of LERK-2-mediated cell adhesion, whereby inflammation is reduced.

The invention pertains to a method of antitumor therapy, comprising administering to a mammal a therapeutically effective amount of an inhibitor of LERK-2-mediated cell adhesion, whereby angiogenesis is reduced, thereby inhibiting tumor growth.

The invention also relates to a method for enhancing angiogenesis, comprising administering to a mammal a therapeutically effective amount of a promoter of LERK-2-mediated cell adhesion, whereby angiogenesis is enhanced.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A-1C show the results of FACS analysis of the expression of LERK-2 (as assessed by binding of anti-LERK-2 antibody 4A1) on leukocytes. Figure 1A shows staining of CD4 positive mononuclear cells, the first column gating on all lymphocytes and the second gating on monocytes; both lymphocytes and monocytes express CD4). Both monocytes and lymphocytes show positive results with anti-LERK-2 antibody 4A1. Figure 1B shows staining using CD14, a monocyte-specific marker, and 4A1. In the first column all cells are largely confined to the lower axis

-5-

while in the second column cells are in the upper right column indicating they are positive for both LERK-2 and CD14 (LERK-2 positive monocytes). Figure 1C shows that LERK-2 reactivity is associated with the memory marker CD45RO. For lymphocytes, the lower right quadrant shows similar levels of reactivity, indicating that CD45RO negative populations (naive cells) non-specifically bind antibody. Cells are seen only in the upper right quadrant for the LERK-2 monoclonal antibody and not the control, indicating that a memory subset of lymphocytes is positive for LERK-2.

Figures 2A and 2B show the results of FACS analysis of the expression of LERK-2 on leukocyte subsets using 4A1 as a staining reagent. Myelomonocytic cell lines including THP-1 and the T cell lines HuT 78 and HSB-2 are brightly positive for LERK-2. Some B cell lines express lower levels of LERK-2, including CA, RAMOS and RAJI. In Figure 2A, LERK-2 expression is also seen in unstimulated HUVEC cultures while expression is slightly increased by 24-hour incubation with TNF- $\alpha$ . The boe tracing represents a control monoclonal antibody (IgG1), while the open tracing is reactivity with anti-LERK-2 monoclonal antibody 4A1.

Figure 3 shows the results of PCR analysis of various cell lines for expression of LERK-2. Results indicate that cells which are positive for LERK-2 expression by FACS analysis can also specifically amplify a 600 base pair DNA fragment by RNA PCR with LERK-2-specific primers. Positive results are seen in lanes 1 (L1-2 LERK-2 transfectants), 3 (peripheral blood mononuclear cells), 4 (HSB-2), 7 (K562), 9 (RAJI), 10 (RAMOS) and 12 (THP-1). Intensity of the signal closely parallels the intensity of the staining observed by FACS analysis.



-6-

Figure 4 show the results of FACS analysis using a LERK-2 Ig chimera. These results indicate that TK1 cells express a receptor for LERK-2.

Figure 5 shows the results of FACS analysis using a LERK-2 Ig chimera. These results indicate that HUVECS also express a receptor for LERK-2.

#### DETAILED DESCRIPTION OF THE INVENTION

LERK-2 (Ligand for *epk*-related kinase-2; Beckmann et al., *EMBO J.* 13(16):3757-3762 (1994)) has been identified as a ligand for the elk and hek tyrosine kinase receptors, which are members of the Eph RPTK family. LERK-2 is a transmembrane protein which is related by sequence homology to LERK-1 (also known as B61; Beckmann et al., *EMBO J.* 13(16):3757-3762 (1994)).

Eph RPTKs are characterized as encoding a structurally-related cysteine rich extracellular domain containing a single immunoglobulin-like loop near the N-terminus and two fibronectin III repeats adjacent to the plasma membrane. The structure of the extracellular region is thought to determine ligand binding specificity. Examples of Eph RPTK family members include mouse Nuk (GenBank Accession No. L25890) and its homologs Hek5 and Cek5 in chickens (Pasquale, *Cell Regulation* 2:523-534 (1991)), Sek3 in mice, Tyro5 in rats (Lai and Lemke, *Neuron* 6:691-704 (1991)) and Erk in humans. Expression of Nuk was found to be essential for formation of at least one commissure in the brain, the medial tract of the anterior commissure, as well as for other brain structures (WO 97/14966 (Pawson et al.)).

As described herein below, an expression cloning system was used to isolate a cDNA from a human mesenteric lymph node library which, upon transfection into CHO/P

-7-

cells, mediates binding of the murine T cell lymphoma, TK1, through the Nuk receptor. Sequencing of the isolated cDNA revealed it to be LERK-2. The interaction of lymphoid cells with LERK-2 represents a step in cell adhesion cascades that has not previously been described.

As described herein, it has been discovered that LERK-2 can mediate cell-cell adhesion between cells expressing LERK-2 and cells expressing a receptor for LERK-2, e.g., cells expressing an Eph RPTK such as Nuk or its homologs. It has further been discovered that Nuk, an RPTK which binds LERK-2, is present on T cell subsets, e.g., TK-1 cell lines, and that LERK-2/Nuk interactions mediate adhesion between cells expressing the ligand and cells expressing the receptor. It has also been discovered that agents which alter the LERK-2/receptor interaction can modulate this cell adhesion, thereby modulating the signals propagated by the stimulation of the receptor through binding of LERK-2. Accordingly, the cellular processes which are regulated by the signals transduced by the LERK-2/RPTK interaction, such as angiogenesis, inflammation and leukocyte trafficking, can be modulated as described herein by modulating LERK-2-mediated cell adhesion.

Angiogenesis, the generation of capillaries, is virtually absent in healthy adult organisms and is restricted to a few instances, including wound healing and the formation of the corpus luteum, endometrium and placenta. Endothelial cells play a key role in the formation of new capillaries, and the neovascularization process occurs via a series of sequential steps, which are similar regardless of the nature of the inducing stimulus. Under normal physiological conditions in the healthy adult mammal, except during cyclical changes in the female

-8-

reproductive tract or in response to wound healing, the coordinated sequential cellular events leading to new capillaries are spatially and temporally restricted so that inhibitory influences on neovascularization  
5 predominate.

However, in certain pathological conditions angiogenesis is dramatically enhanced and is no longer self-limited (Maugh, *Science* 212:1374-1375 (1981); Auerback, *Lymphokines* 4:69 (1981)). Angiogenesis is a  
10 major component of some opthalmological pathologies such as corneal graft rejection, corneal neovascularization following injury or infection, diabetic retinopathy, retrolental fibroplasia and neovascular glaucoma (Garner, *Int. Rev. Exp. Pathol.* 284:249-307 (1986)). It is also a  
15 significant factor in many ulcerative diseases such as rheumatoid arthritis, ulcerative colitis and gastric ulcer. In addition, angiogenesis is a major component in pathological but nonmalignant conditions such as hemangioma, angiofibroma of the nasopharynx, avascular  
20 necrosis of bone, and psoriasis.

Perhaps the clinically most important manifestation of pathological angiogenesis is that induced by solid tumors (Folkman, *Adv. Cancer Res.* 43:175-203 (1985); Folkman, *Nature Medicine* 1(1) (1995)). In order for a  
25 neoplasm to grow progressively as a solid mass consisting of layers of living cells more than a few millimeters thick, it must induce nearby capillaries to sprout and develop a new vascular network around and within the tumor. The new vascular network supplies the tumor with  
30 vital nutrients and oxygen and provides a removal route for toxic products of active cell metabolism. Furthermore, new tumor vessels provide an exit for tumor cells to metastize to distant sites. Thus, the

-9-

progressive growth of a solid tumor to develop into a life-threatening malignancy is strictly dependent on angiogenesis.

Members of the LERK ligand family have been shown to be involved in the angiogenic process. For example, LERK-1 has been shown in vitro to be angiogenic for human umbilical vein endothelial cells (HUVECs) via the Eck receptor, and has been shown to promote corneal neovascularization in a rat model (Pandey et al., *Science* 268:567-569 (1995)). LERK-2 has been shown to be angiogenic for a human renal microvascular endothelial cell line via the Elk receptor (HRMEC; Daniel et al., *Kidney International* 50(Suppl. 57):S-73-S-81 (1996)).

LERK-1 has also been shown to be an immediate early response gene in endothelial cells; its expression is induced by proinflammatory factors such as tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), interleukin-1 and lipopolysaccharide (Holzman et al., *Mol. Cell. Biol.* 10:5830-5838 (1990)). LERK-2 has similarly been shown to be induced by TNF- $\alpha$  in endothelial cells (Beckmann et al., *EMBO J.* 13:3757-3762 (1994)).

Inflammatory disorders or conditions generally share several features which are agreed to be characteristic of the inflammation process. These include fenestration of the microvasculature, leakage of blood elements into the interstitial spaces, and migration of leukocytes into the inflamed tissue. On a macroscopic level, this is usually accompanied by the familiar clinical signs of erythema, edema, tenderness (hyperalgesia), and pain. Inflammatory diseases such as chronic inflammatory pathologies and vascular inflammatory pathologies include, for example, aneurysms, hemorrhoids, sarcoidosis, chronic inflammatory bowel disease, delayed type hypersensitivity reaction,

-10-

allograft rejection, ulcerative colitis, Crohn's disease, disseminated intravascular coagulation, atherosclerosis, and Kawasaki's pathology. Inflammatory diseases also include coronary artery disease, hypertension, stroke, 5 asthma, psoriasis, nephritis, chronic hepatitis, multiple sclerosis, peripheral neuropathy, chronic or recurrent sore throat, laryngitis, tracheobronchitis, chronic vascular headaches (including migraines, cluster headaches and tension headaches), pneumonia and bacterial or viral 10 infection.

The present invention relates to a method of modulating (i.e., inhibiting or promoting) LERK-2-mediated cell adhesion. In a preferred embodiment, the invention pertains to a method of modulating LERK-2/Nuk-mediated 15 cell adhesion (i.e., cell adhesion mediated by the interaction or binding of LERK-2 and Nuk). As used herein, "LERK-2-mediated cell adhesion" means cell adhesion mediated by binding of LERK-2 to its corresponding Eph RPTK (i.e., a LERK-2 receptor or 20 receptor for LERK-2), thereby mediating adhesion between the cell which expresses LERK-2 and the cell which expresses the Eph RPTK. Eph RPTKs which bind to LERK-2 include, but are not limited to, Cek10, Sek-4, Tyro6, HEK2, Cek5, Nuk (Sek-3), Tyro5, ERK, Cek6 and Elk. The 25 method of the invention comprises contacting at least one cell selected from the group consisting of a cell expressing LERK-2 and a cell expressing a receptor for LERK-2 with an inhibitor or promoter of LERK-2-mediated cell adhesion. Suitable cells can be cells which normally 30 (in nature) express LERK-2 or a receptor for LERK-2, or can be cells which have been engineered, e.g., by recombinant technology, to express LERK-2 or a receptor for LERK-2. *In vivo* expression can be accomplished via

-11-

somatic cell expression according to suitable methods (see, e.g. U.S. Patent No. 5,399,346). In this embodiment, nucleic acid encoding the protein can be incorporated into a retroviral, adenoviral or other  
5 suitable vector (preferably, a replication deficient infectious vector) for delivery, or can be introduced into a transfected or transformed host cell capable of expressing the protein for delivery.

In particular embodiments, the cells expressing LERK-  
10 2 or receptor for LERK-2 are lymphoid cells (e.g., leukocytes, including lymphocytes such as T cells) or endothelial cells. In a preferred embodiment, the receptor for LERK-2 is selected from the group consisting of Nuk, Cek5, Tyro5 and ERK. In a particularly preferred  
15 embodiment the LERK-2 receptor is Nuk.

Suitable inhibitors for use in the invention include, but are not limited to, anti-LERK-2 antibodies or functional antibody fragments thereof, anti-Eph RPTK antibodies or functional antibody fragments thereof, e.g.,  
20 anti-Nuk antibodies, soluble LERK-2 and LERK-2-Ig chimeras. Suitable promoters include isolated LERK-2 or LERK-2 chimeras, as well as LERK-2 mimics. As used herein, a "LERK-2 mimic" is an agent which can activate, e.g., by binding, a LERK-2 receptor, e.g., an Eph RPTK  
25 such as Nuk, to induce the same or similar downstream signaling events as those induced by LERK-2. LERK-2 mimics can include, for example, peptides, small molecules and antibodies which can activate a LERK-2 receptor.

For instance, polyclonal and monoclonal antibodies  
30 which bind to LERK-2 or a receptor for LERK-2 are within the scope of the invention. A mammal, such as a mouse, hamster or rabbit, can be immunized with an immunogenic form of LERK-2 or LERK-2 receptor (i.e., an antigenic

-12-

fragment of LERK-2 or LERK-2 receptor which is capable of eliciting an antibody response) such as isolated and/or recombinant peptide (including synthetic molecules, such as synthetic peptides). Preparation of immunizing antigen  
5 (e.g., a peptide conjugated to a suitable carrier, such as keyhole limpet hemocyanin, serum albumin, etc.), and polyclonal and monoclonal antibody production can be performed using any suitable technique. Techniques for conferring immunogenicity on a protein or peptide include  
10 conjugation to carriers or other techniques well known in the art. The protein or polypeptide can be administered in the presence of an adjuvant. The progress of immunization can be monitored by detection of antibody titers in plasma or serum. Standard ELISA or other  
15 immunoassays can be used with the immunogen as antigen to assess the levels of antibody.

Following immunization, anti-peptide antisera can be obtained, and if desired, polyclonal antibodies can be isolated from the serum. Monoclonal antibodies can also  
20 be produced by standard techniques which are well known in the art (see e.g., Kohler et al., *Nature*, 256: 495-497 (1975) and *Eur. J. Immunol.* 6: 511-519 (1976); Milstein et al., *Nature* 266: 550-552 (1977); Koprowski et al., U.S. Patent No. 4,172,124; Harlow, E. and D. Lane, 1988,  
25 *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory: Cold Spring Harbor, NY); *Current Protocols In Molecular Biology*, Vol. 2 (Supplement 27, Summer '94), Ausubel, F.M. et al., Eds., (John Wiley & Sons: New York, NY), Chapter 11, (1991)).

30 One standard approach to the production of monoclonal antibodies is that described by Kohler et al., (1975). In this method a subject animal (e.g., mouse, rat or other rodent) is immunized with the material against which the

-13-

antibodies are desired (e.g., LERK-2 or portion thereof or LERK-2 receptor or portion thereof). The animal's immune system generates antibodies against the material via its B cells, which are localized in the animal's spleen. The spleen (or lymph) is removed and treated to isolate individual cells. These cells are fused with cell lines which are immortal in culture (e.g., a myeloma cell line such as SP2/0) in the presence of an agent which facilitates cell fusion (e.g., polyethylene glycol). The fused cells yield hybridomas, each of which produces antibodies of a particular epitopic specificity. Suitable screening procedures are then carried out to identify and isolate hybridomas producing antibodies with the desired epitopic specificity (monoclonal antibodies). For example, cells which produce antibodies with the desired specificity can be selected by a suitable assay (e.g., ELISA).

In order to identify the particular epitopic specificity of a selected monoclonal antibody, a panel of chimeras comprising selected portions of the antigen in combination with selected portions of another suitable molecule can be constructed by transferring restriction fragments flanked by specific restriction sites (e.g., BamHI, AflII, ClaI, EcoRI, and XbaI sites). The constructs can be transferred into an appropriate vector, under control of an appropriate promoter, and transfected in CHO-k1 cells as described by Perret et al. (*Biochem. Biophys. Res. Commun.* 17:1044 (1990)). Transfectants can be stained with the selected monoclonal antibody to determine the epitopic specificity of the particular antibody.

Monoclonal antibodies 2A1 and 4A1 were deposited under the terms of the Budapest Treaty at the American



-14-

Type Culture Collection, 10801 University Boulevard,  
Manassas, Virginia 20110, on behalf of LeukoSite, Inc.,  
215 First Street, Cambridge, MA 02142, on October 2, 1997,  
and were assigned ATCC Accession Nos. HB-12412 and HB-  
5 12413, respectively.

In a preferred embodiment of the invention, the  
inhibitor is selected from the group consisting of  
monoclonal antibody 2A1 (ATCC Accession No. HB-12412),  
monoclonal antibody 4A1 (ATCC Accession No. HB-12413),  
10 functional antibody fragments of 2A1, functional antibody  
fragments of 4A1, an antibody or functional antibody  
fragment thereof which can compete with monoclonal  
antibody 2A1 or 4A1 for binding to LERK-2 or portion  
thereof, or an antibody or functional antibody fragment  
15 thereof which has the same or similar epitopic specificity  
as that of 2A1 or 4A1. In a particularly preferred  
embodiment, the inhibitor (e.g., antibody or functional  
antibody fragment) selectively inhibits the interaction of  
LERK-2 with Nuk.

20 The invention also relates to monoclonal antibody  
2A1, monoclonal antibody 4A1, functional antibody  
fragments of 2A1, functional antibody fragments of 4A1, an  
antibody or functional antibody fragment thereof which can  
compete with monoclonal antibody 2A1 or 4A1 for binding to  
25 LERK-2 or portion thereof, or an antibody or functional  
antibody fragment thereof which has the same or similar  
epitopic specificity as that of 2A1 or 4A1. Also included  
within the scope of the invention are the hybridoma cell  
line deposited under ATCC Accession No. HB-12412, the  
30 hybridoma cell line deposited under ATCC Accession No. HB-  
12413, a monoclonal antibody or antigen binding fragment  
thereof produced by the hybridoma cell line deposited  
under ATCC Accession No. HB-12412, and a monoclonal

-15-

antibody or antigen binding fragment thereof produced by the hybridoma cell line deposited under ATCC Accession No. HB-12413. These antibodies have utility as inhibitors of LERK-2-mediated cell adhesion, as well as reagents for immunostaining of cells and tissues to identify expression of LERK-2.

The cell lines of the present invention have uses other than for the production of the monoclonal antibodies. For example, the cell lines of the present invention can be fused with other cells (such as suitably drug-marked human myeloma, mouse myeloma, human-mouse heteromyeloma or human lymphoblastoid cells) to produce additional hybridomas, and thus provide for the transfer of the genes encoding the monoclonal antibodies. In addition, the cell lines can be used as a source of nucleic acids encoding the anti-LERK-2 immunoglobulin chains, which can be isolated and expressed (e.g., upon transfer to other cells using any suitable technique (see e.g., Cabilly et al., U.S. Patent No. 4,816,567; Winter, U.S. Patent No. 5,225,539)).

Other suitable methods of producing or isolating antibodies which inhibit LERK-2-mediated cell adhesion can be used, including, for example, methods which select recombinant antibody from a library, or which rely upon immunization of transgenic animals (e.g., mice) capable of producing a full repertoire of human antibodies (see e.g., Jakobovits et al., *Proc. Natl. Acad. Sci. USA*, 90: 2551-2555 (1993); Jakobovits et al., *Nature*, 362: 255-258 (1993); Lonberg et al., U.S. Patent No. 5,545,806; Surani et al., U.S. Patent No. 5,545,807).

Single chain antibodies, and chimeric, humanized or primatized (CDR-grafted) antibodies, as well as chimeric or CDR-grafted single chain antibodies, comprising

-16-

portions derived from different species, are also encompassed by the present invention and the term "antibody". The various portions of these antibodies can be joined together chemically by conventional techniques, or can be prepared as a contiguous protein using genetic engineering techniques. For example, nucleic acids encoding a chimeric or humanized chain can be expressed to produce a contiguous protein. See, e.g., Cabilly et al., U.S. Patent No. 4,816,567; Cabilly et al., *European Patent* 10 *No. 0,125,023 B1*; Boss et al., U.S. Patent No. 4,816,397; Boss et al., *European Patent No. 0,120,694 B1*; Neuberger, M.S. et al., WO 86/01533; Neuberger, M.S. et al., *European Patent No. 0,194,276 B1*; Winter, U.S. Patent No. 5,225,539; and Winter, *European Patent No. 0,239,400 B1*. See also, Newman, R. et al., *BioTechnology*, 10: 1455-1460 (1992), regarding primatized antibody, and Ladner et al., U.S. Patent No. 4,946,778 and Bird, R.E. et al., *Science*, 242: 423-426 (1988) regarding single chain antibodies.

In addition, functional fragments of antibodies, including fragments of chimeric, humanized, primatized or single chain antibodies, can also be produced. Functional fragments of the foregoing antibodies retain at least one modulation function of the full-length antibody from which they are derived. Preferred functional fragments retain an antigen binding function of a corresponding full-length antibody (e.g., specificity for LERK-2 or an Eph receptor for LERK-2). Particularly preferred functional fragments retain the ability to inhibit LERK-2-mediated cell adhesion. For example, in one embodiment, a functional 25 fragment can inhibit the interaction of LERK-2 with a receptor for LERK-2 (e.g., Nuk), thereby inhibiting one or more receptor-mediated functions, such as angiogenesis, inflammation, or leukocyte trafficking.

-17-

For example, antibody fragments capable of binding to LERK-2 or portion thereof or to a LERK-2 receptor or portion thereof, including, but not limited to, Fv, Fab, Fab' and F(ab')<sub>2</sub> fragments, are encompassed by the invention. Such fragments can be produced by enzymatic cleavage or by recombinant techniques. For instance, papain or pepsin cleavage can generate Fab or F(ab')<sub>2</sub> fragments, respectively. Antibodies can also be produced in a variety of truncated forms using antibody genes in which one or more stop codons has been introduced upstream of the natural stop site. For example, a chimeric gene encoding a F(ab')<sub>2</sub> heavy chain portion can be designed to include DNA sequences encoding the CH<sub>1</sub> domain and hinge region of the heavy chain.

The term "humanized immunoglobulin" as used herein refers to an immunoglobulin comprising portions of immunoglobulins of different origin, wherein at least one portion is of human origin. Accordingly, the present invention relates to a humanized immunoglobulin having binding specificity for LERK-2 or a receptor for LERK-2, said immunoglobulin comprising an antigen binding region of nonhuman origin (e.g., rodent) and at least a portion of an immunoglobulin of human origin (e.g., a human framework region, a human constant region or portion thereof). For example, the humanized antibody can comprise portions derived from an immunoglobulin of nonhuman origin, such as a mouse, and from immunoglobulin sequences of human origin (e.g., a chimeric immunoglobulin), joined together chemically by conventional techniques (e.g., synthetic) or prepared as a contiguous polypeptide using genetic engineering techniques (e.g., DNA encoding the protein portions of the chimeric antibody can be expressed to produce a contiguous

-18-

polypeptide chain). Another example of a humanized immunoglobulin of the present invention is an immunoglobulin containing one or more immunoglobulin chains comprising a CDR of nonhuman origin (e.g., one or  
5 more CDRs derived from an antibody of nonhuman origin) and a framework region derived from a light and/or heavy chain of human origin (e.g., CDR-grafted antibodies with or without framework changes). In one embodiment, the humanized immunoglobulin can compete with 2A1 or 4A1  
10 monoclonal antibody for binding to LERK-2. Chimeric or CDR-grafted single chain antibodies are also encompassed by the term humanized immunoglobulin. See, e.g., Cabilly et al., U.S. Patent No. 4,816,567; Cabilly et al., European Patent No. 0,125,023 B1; Queen et al., European  
15 Patent No. 0,451,216 B1; Boss et al., U.S. Patent No. 4,816,397; Boss et al., European Patent No. 0,120,694 B1; Neuberger, M.S. et al., WO 86/01533; Neuberger, M.S. et al., European Patent No. 0,194,276 B1; Winter, U.S. Patent No. 5,225,539; Winter, European Patent No. 0,239,400 B1;  
20 Padlan, E.A. et al., European Patent Application No. 0,519,596 A1. See also, Ladner et al., U.S. Patent No. 4,946,778; Huston, U.S. Patent No. 5,476,786; and Bird, R.E. et al., *Science*, 242: 423-426 (1988)), regarding single chain antibodies.

25 Such humanized immunoglobulins can be produced using synthetic and/or recombinant nucleic acids to prepare genes (e.g., cDNA) encoding the desired humanized chain. For example, nucleic acid (e.g., DNA) sequences coding for humanized variable regions can be constructed using PCR  
30 mutagenesis methods to alter DNA sequences encoding a human or humanized chain, such as a DNA template from a previously humanized variable region (see e.g., Kamman, M., et al., *Nucl. Acids Res.*, 17: 5404 (1989)); Sato, K.,

-19-

et al., *Cancer Research*, 53: 851-856 (1993); Daugherty, B.L. et al., *Nucleic Acids Res.*, 19(9): 2471-2476 (1991); and Lewis, A.P. and J.S. Crowe, *Gene*, 101: 297-302 (1991)). Using these or other suitable methods, variants  
5 can also be readily produced. In one embodiment, cloned variable regions can be mutagenized, and sequences encoding variants with the desired specificity can be selected (e.g., from a phage library; see e.g., Krebber et al., U.S. 5,514,548; Hoogenboom et al., WO 93/06213,  
10 published April 1, 1993)).

Other appropriate inhibitors and promoters can be identified in a suitable assay. For example, the invention pertains to a method of detecting or identifying an inhibitor or promoter of LERK-2-mediated cell adhesion,  
15 comprising combining an agent to be tested with a composition comprising a cell expressing LERK-2 and a composition comprising a cell expressing a receptor for LERK-2 (e.g., Nuk), under conditions suitable for binding of LERK-2 to the receptor for LERK-2. The cell adhesion  
20 between the cell expressing LERK-2 and the cell expressing a receptor for LERK-2 is detected or measured, and inhibition or promotion of cell adhesion by the agent is indicative that the agent is an inhibitor or promoter, respectively.

25 In a particular embodiment, cell adhesion between a cell expressing LERK-2 and a cell expressing a receptor for LERK-2 mediates a cellular signalling and/or cellular response (e.g., angiogenesis, inflammatory response, leukocyte migration), and cell adhesion is monitored by  
30 detecting or measuring a signalling activity or cellular response of one or both of the cells in response thereto. Signalling events triggered by LERK-2/receptor interaction and LERK-2-mediated cell adhesion, such as induction of

-20-

rapid and transient increase in the concentration of intracellular (cytosolic) free calcium  $[Ca^{2+}]_i$ , can be assayed by methods known in the art or other suitable methods (see e.g., Neote, K. et al., *Cell*, 72: 415-425 1993); Van Riper et al., *J. Exp. Med.*, 177: 851-856 (1993); Dahinden, C.A. et al., *J. Exp. Med.*, 179: 751-756 (1994)).

For example, modulators which can be identified as described herein include modulating antibodies, e.g., antibodies which inhibit or promote LERK-2-mediated cell adhesion. Modulating antibodies can be an antibody or functional antibody fragment which modulates LERK-2-mediated cell adhesion, such as an antibody or fragment which modulates binding of LERK-2 to a receptor for LERK-2. In a particular embodiment, the antibody or functional antibody fragment is an anti-LERK-2 antibody.

Binding inhibition assays can also be used to identify antibodies which bind LERK-2 or a receptor for LERK-2 and inhibit binding of another compound such as a receptor for LERK-2 or LERK-2, respectively, thereby inhibiting LERK-2-mediated cell adhesion. For example, a binding assay can be conducted in which a reduction in the binding of LERK-2 to a LERK-2 receptor (in the absence of an antibody), as compared to binding of LERK-2 to a LERK-2 receptor in the presence of the antibody, is detected or measured. A reduction in the extent of binding in the presence of the antibody is indicative of inhibition of binding by the antibody.

In one embodiment, direct inhibition of the binding of LERK-2 to a receptor for LERK-2 by an inhibitor, e.g., an antibody, is monitored. For example, the ability of an antibody to inhibit the binding of  $^{125}I$ -labeled LERK-2 to Nuk can be monitored. Such an assay can be conducted

-21-

using either whole cells (e.g., T cells, or a suitable cell line containing nucleic acid encoding the ligand or receptor) or a membrane fraction from said cells, for instance.

5           It will be understood that the inhibitory effect of antibodies of the present invention can be assessed in a binding inhibition assay. Competition between antibodies for receptor and/or ligand (LERK-2) binding can also be assessed in the method. Antibodies which are identified  
10 in this manner can be further assessed to determine whether, subsequent to binding, they act to inhibit LERK-2-mediated cell adhesion and/or to assess their therapeutic utility.

Chemotaxis assays can also be used to assess the  
15 ability of a compound to block binding of LERK-2 to a LERK-2 receptor and/or inhibit LERK-2-mediated cell adhesion. These assays are based on the functional migration of cells *in vitro* or *in vivo* induced by a compound. The use of an *in vitro* transendothelial  
20 chemotaxis assay is described by Springer et al. (Springer et al., WO 94/20142, published September 15, 1994, the teachings of which are incorporated herein by reference; see also Berman et al., *Immunol. Invest.* 17: 625-677 (1988)). Migration across endothelium into collagen gels  
25 has also been described (Kavanaugh et al., *J. Immunol.*, 146: 4149-4156 (1991)). Stable transfectants of mouse L1-2 pre-B cells or of other suitable host cells capable of chemotaxis can be used in chemotaxis assays, for example.

30           Generally, chemotaxis assays monitor the directional movement or migration of a suitable cell (such as a leukocyte (e.g., lymphocyte, eosinophil, basophil)) into or through a barrier (e.g., endothelium, a filter), toward



-22-

increased levels of a compound, from a first surface of the barrier toward an opposite second surface. Membranes or filters provide convenient barriers, such that the directional movement or migration of a suitable cell into or through a filter, toward increased levels of a compound, from a first surface of the filter toward an opposite second surface of the filter, is monitored. In some assays, the membrane is coated with a substance to facilitate adhesion, such as ICAM-1, fibronectin or collagen. Such assays provide an *in vitro* approximation of leukocyte "homing".

For example, one can detect or measure inhibition of the migration of cells in a suitable container (a containing means), from a first chamber into or through a microporous membrane into a second chamber which contains an antibody to be tested, and which is divided from the first chamber by the membrane. A suitable membrane, having a suitable pore size for monitoring specific migration in response to compound, including, for example, nitrocellulose, polycarbonate, is selected. For example, pore sizes of about 3-8 microns, and preferably about 5-8 microns can be used. Pore size can be uniform on a filter or within a range of suitable pore sizes.

To assess migration and inhibition of migration, the distance of migration into the filter, the number of cells crossing the filter that remain adherent to the second surface of the filter, and/or the number of cells that accumulate in the second chamber can be determined using standard techniques (e.g., microscopy). In one embodiment, the cells are labeled with a detectable label (e.g., radioisotope, fluorescent label, antigen or epitope label), and migration can be assessed in the presence and absence of the antibody by determining the presence of the

-23-

label adherent to the membrane and/or present in the second chamber using an appropriate method (e.g., by detecting radioactivity, fluorescence, immunoassay). The extent of migration induced by an antibody agonist can be  
5 determined relative to a suitable control (e.g., compared to background migration determined in the absence of the antibody, compared to the extent of migration induced by a second compound (i.e., a standard), compared with migration of untransfected cells induced by the antibody).

10 In a preferred embodiment, particularly for T cells, monocytes or other cells expressing LERK-2, transendothelial migration is monitored. Such assays are better physiological models, because they more accurately recapitulate *in vivo* conditions in which leukocytes  
15 emigrate from blood vessels toward chemoattractants present in the tissues at sites of inflammation by crossing the endothelial cell layer lining the vessel wall. In addition, transendothelial assays have lower background and as a result a higher signal to noise ratio.

20 In one embodiment used to test for an antibody inhibitor, a composition comprising cells capable of migration and expressing a receptor for LERK-2 can be placed in the first chamber. A composition comprising LERK-2 or a promoter capable of inducing chemotaxis of the  
25 cells in the first chamber (having chemoattractant function) is placed in the second chamber. Preferably shortly before the cells are placed in the first chamber, or simultaneously with the cells, a composition comprising the antibody to be tested is placed, preferably, in the  
30 first chamber. Antibodies which can bind receptor and inhibit the induction of chemotaxis, by LERK-2 or promoter, of the cells expressing a receptor for LERK-2 in this assay are inhibitors of receptor function. A

-24-

reduction in the extent of migration induced by LERK-2 or promoter in the presence of the antibody is indicative of inhibitory activity. Separate binding studies can be performed to determine whether inhibition is a result of binding of the antibody to receptor or occurs via a different mechanism.

In vivo assays which monitor leukocyte infiltration of a tissue, in response to injection of a compound (e.g., antibody) in the tissue, are models of *in vivo* homing which measure the ability of cells to respond to a ligand or promoter by emigration and chemotaxis to a site of inflammation.

The assays described above, which can be used to assess the inhibitory or promoter effects of test compounds, can be adapted to identify inhibitors and/or promoters of LERK-2-mediated cell adhesion. For example, agents having the same or a similar binding specificity as that of an antibody of the present invention or functional portion thereof can be identified by a competition assay with said antibody or portion thereof. In one embodiment, cells expressing LERK-2 or functional variant thereof (e.g., leukocytes or suitable host cells which have been engineered to express LERK-2 or functional variant encoded by a nucleic acid introduced into said cells) are used in an assay to identify and assess the efficacy of substances which bind LERK-2 or LERK-2 receptors, including inhibitors or promoters of receptor function.

The present invention also pertains to a method for the treatment of angiogenic disease, particularly angiogenesis-sustained solid tumors, in an individual. The term "individual" is defined herein to include animals such as mammals, including, but not limited to, primates (e.g., humans), cows, sheep, goats, horses, dogs, cats,

-25-

rabbits, guinea pigs, rats, mice or other bovine, ovine, equine, canine, feline, rodent or murine species. As used herein, "angiogenic disease" includes any pathological condition associated with enhanced angiogenesis, that is, a condition which is directly or indirectly supported, sustained or aggravated by increased angiogenesis.

As all solid tumors are dependent on enhanced angiogenesis to grow, the treatment of the invention is generally applicable to any solid tumor; the invention is also useful in the treatment or prevention of metastasis. For example, the methods of the present invention are useful in the treatment of tumors of the prostate, breast, endometrium, lung, colon and kidney, sarcomas, melanomas and skin tumors, including metastasizing tumors. The invention also relates to treatments for psoriasis, rheumatoid arthritis, atherosclerosis, ulcerative colitis, gastric ulcers, corneal graft rejection, corneal neovascularization following injury or infection, diabetic retinopathy, retrolental fibroplasia and neovascular glaucoma, as well as conditions such as hemangioma, angiofibroma of the nasopharynx, avascular necrosis of bone, and psoriasis. The method of the present invention can also be used to inhibit the angiogenesis which accompanies the development of the endometrium, thereby finding use as a contraceptive. In some instances it may be clinically beneficial to enhance or promote angiogenesis in an individual. For example, the process of wound healing requires the neovascularization of newly generated tissue, and a major factor in graft survival is the ability of new vasculature to infiltrate the grafted tissue and connect the tissue to the nutrients and waste removal system provided by the circulatory system of the individual. Enhanced angiogenesis is also useful in the

-26-

treatment of frostbite to replace the damaged or destroyed capillaries in frostbitten tissue, as well as in recannulization of a thrombus following vascular thrombosis.

5           Accordingly, the invention pertains to a method of modulating (i.e., inhibiting or promoting) angiogenesis in an individual in need of such therapy, comprising administering a therapeutically effective amount of an agent which inhibits (an inhibitor) or promotes (a  
10   promoter) LERK-2-mediated cell adhesion to an individual in need of such therapy. A therapeutically effective amount is an amount sufficient to achieve the desired therapeutic effect, under the conditions of administration, such as an amount sufficient for  
15   inhibition or promotion of LERK-2-mediated cell adhesion, and thereby, inhibition or promotion, respectively, of a LERK-2/receptor-mediated process (e.g., an angiogenic response, an inflammatory response or a leukocyte trafficking response).

20           In one embodiment, a compound which inhibits LERK-2-mediated cell adhesion is administered to inhibit angiogenesis. As used herein, "inhibition" includes any quantitative or qualitative reduction, including prevention and complete absence, relative to the control.

25   In another embodiment, a compound which promotes LERK-2-mediated cell adhesion is administered to promote angiogenesis. As used herein, "promotion" includes any quantitative or qualitative increase relative to the control. For example, the invention pertains to a method  
30   of modulating angiogenesis in an individual in need of such therapy, comprising administering an agent which inhibits or promotes LERK-2/Nuk-mediated cell adhesion.

-27-

Additionally, the invention relates to a method for modulating the migration of cells, particularly endothelial cells. Cell migration is important to angiogenesis. New capillaries cannot be formed unless the  
5 endothelial cells have the ability to migrate to and in the extracellular space. There, they align to form solid cords; formation of a lumen eventually occurs when cords sprouting from adjacent capillaries meet each other.

In another aspect, the present invention provides a  
10 method of modulating (i.e., inhibiting or promoting) an inflammatory response in an individual in need of such therapy, comprising administering a therapeutically effective amount of an agent which inhibits or promotes LERK-2-mediated cell adhesion to an individual in need of  
15 such therapy. For example, the invention pertains to a method of modulating inflammation in an individual in need of such therapy, comprising administering an agent which inhibits or promotes LERK-2/Nuk-mediated cell adhesion. In one embodiment, a compound which inhibits LERK-2-  
20 mediated cell adhesion is administered to inhibit inflammation. As a result, one or more inflammatory processes, such as leukocyte emigration, chemotaxis, exocytosis (e.g., of enzymes) or inflammatory mediator release, is inhibited. For example, leukocytic  
25 infiltration of inflammatory sites (e.g., in a delayed-type hypersensitivity response) can be inhibited according to the present method.

In another embodiment, a compound which promotes LERK-2-mediated cell adhesion is administered to promote  
30 an inflammatory response, such as leukocyte emigration, chemotaxis, exocytosis (e.g., of enzymes) or inflammatory mediator release, resulting in the beneficial stimulation of inflammatory processes. For example, natural killer

-28-

cells can be recruited to combat viral infections or neoplastic disease.

A variety of *in vivo* models of inflammation are available, which can be used to assess the effects of inhibitors or promoters of LERK-2-mediated cell adhesion *in vivo* as therapeutic agents, including a sheep model for asthma (see e.g., Weg, V.B. et al., *J. Exp. Med.*, 177: 561 (1993)), a rat delayed type hypersensitivity model (Rand, M.L. et al., *Am. J. Pathol.*, 148: 855-864 (1996)), or other suitable models.

Inflammatory diseases or conditions, including chronic diseases, of humans or other species which can be treated with inhibitors of LERK-2-mediated cell adhesion, include, but are not limited to allergic diseases and conditions, including systemic anaphylaxis or hypersensitivity responses, drug allergies (e.g., to penicillin, cephalosporins), insect sting allergies; inflammatory bowel diseases, such as Crohn's disease, ulcerative colitis, ileitis and enteritis; vaginitis; psoriasis and inflammatory dermatoses such as dermatitis, eczema, atopic dermatitis, allergic contact dermatitis, urticaria; vasculitis (e.g., necrotizing, cutaneous, and hypersensitivity vasculitis); spondyloarthropathies; scleroderma; respiratory allergic diseases such as asthma, allergic rhinitis, hypersensitivity lung diseases, hypersensitivity pneumonitis, and interstitial lung diseases (ILD) (e.g., idiopathic pulmonary fibrosis, or ILD associated with rheumatoid arthritis, or other autoimmune conditions).

Other disease and conditions which can be treated by the methods of this invention include autoimmune diseases, such as arthritis (e.g., rheumatoid arthritis, psoriatic arthritis), multiple sclerosis, systemic lupus

-29-

erythematosus, myasthenia gravis, diabetes, including diabetes mellitus and juvenile onset diabetes, glomerulonephritis and other nephritides, autoimmune thyroiditis, Behcet's disease; graft rejection (e.g., in  
5 transplantation), including allograft rejection or graft-versus-host disease; and other diseases or conditions in which undesirable inflammatory responses are to be inhibited can be treated, including, but not limited to, atherosclerosis, cytokine-induced toxicity, myositis  
10 (including polymyositis, dermatomyositis).

Also suitable for treatment as described herein are cancers, particularly those with leukocytic infiltration of the skin or organs such as cutaneous T cell lymphoma (e.g., mycosis fungoides); diseases in which angiogenesis  
15 or neovascularization plays a role, including neoplastic disease, retinopathy (e.g., diabetic retinopathy), and macular degeneration; and infectious diseases, such as bacterial infections and tuberculoid leprosy, and especially viral infections. Inhibitory compounds  
20 described herein can also be administered to inhibit angiogenesis as an antitumor therapy in conjunction with anti-tumor chemotherapeutic agents known in the art.

Inhibitors and promoters of LERK-2-mediated cell adhesion can be used in the described methods.  
25 Appropriate inhibitors and promoters can be identified in a suitable assay, such as assays described herein, and further assessed for therapeutic effect. For example, antibodies of the present invention, including mAb 2A1 and 4A1, as well as functional antibody fragments of mAb 2A1  
30 and 4A1, and antibodies which can compete with monoclonal antibodies 2A1 and 4A1, can be used in the method to inhibit LERK-2-mediated cell adhesion, thereby inhibiting angiogenesis and/or inflammation.



-30-

The present invention also relates to preparations for use in the modulation of angiogenesis and inflammation, the treatment of angiogenic diseases and inflammatory disorders, and the modulation of leukocyte trafficking, the preparation including an inhibitor or promoter of LERK-2-mediated cell adhesion, together with a physiologically acceptable carrier and optionally other physiologically acceptable adjuvants.

According to the method, a therapeutically effective amount of one or more agents (e.g., a preparation comprising an inhibitor or promoter of LERK-2-mediated cell adhesion, particularly LERK-2/Nuk-mediated cell adhesion, can be administered to an individual by an appropriate route, either alone or in combination with another drug.

A variety of routes of administration are possible including, but not limited to, oral, dietary, topical, parenteral (e.g., intravenous, intraarterial, intramuscular, subcutaneous injection), and inhalation (e.g., intrabronchial, intranasal or oral inhalation, intranasal drops) routes of administration, depending on the agent and disease or condition to be treated. For respiratory allergic diseases such as asthma, inhalation is a preferred mode of administration.

Formulation of an agent to be administered will vary according to the route of administration selected (e.g., solution, emulsion, capsule). An appropriate composition comprising the agent to be administered can be prepared in a physiologically acceptable vehicle or carrier. For solutions or emulsions, suitable carriers include, for example, aqueous or alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles can include sodium chloride solution,

-31-

Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's or fixed oils, for instance. Intravenous vehicles can include various additives, preservatives, or fluid, nutrient or electrolyte replenishers and the like  
5 (See, generally, *Remington's Pharmaceutical Sciences*, 17th Edition, Mack Publishing Co., PA, 1985). For inhalation, the agent can be solubilized and loaded into a suitable dispenser for administration (e.g., an atomizer, nebulizer or pressurized aerosol dispenser).

10 Furthermore, where the agent is a protein or peptide, the agent can be administered via *in vivo* expression of the recombinant protein. *In vivo* expression can be accomplished via somatic cell expression according to suitable methods (see, e.g. U.S. Patent No. 5,399,346).  
15 In this embodiment, nucleic acid encoding the protein can be incorporated into a retroviral, adenoviral or other suitable vector (preferably, a replication deficient infectious vector) for delivery, or can be introduced into a transfected or transformed host cell capable of  
20 expressing the protein for delivery. In the latter embodiment, the cells can be implanted (alone or in a barrier device), injected or otherwise introduced in an amount effective to express the protein in a therapeutically effective amount.

25 The present invention will now be illustrated by the following Examples, which are not intended to be limiting in any way. The teachings of all references cited herein are incorporated herein by reference.

-32-

## EXAMPLES

## METHODOLOGY

## RNA Isolation and Selection of Message

Total RNA was isolated from both primate (macaque)  
5 and human (both histologically normal and inflamed ileal  
nodes from a patient with Chrohn's disease) mesenteric  
lymph nodes or tissue culture cells by use of the cesium  
trifluoroacetate (CsTFA™) reagent from Pharmacia (Catalog  
#17-087-02). Tissue was first snap frozen in liquid  
10 nitrogen and subjected to dounce homogenization in a  
solution consisting of 5.5 M guanidinium thiocyanate, 25  
mM sodium citrate, 0.5% sodium laurel sarcosine and 0.2 M  
2-mercaptoethanol, while tissue culture cells ( $1-5 \times 10^8$ )  
were washed once in phosphate buffered saline (PBS) and  
15 homogenized by pipetting. A clarified lysate was then  
layered on a cushion of CsTFA™ and total RNA was pelleted  
by centrifugation for 20 hours at 30,000 RPM.

mRNA was selected by the polyAtract mRNA isolation  
system from Promega. The system uses a biotinylated  
20 oligo(dT) primer to hybridize (in solution) to poly A  
tails of eukaryotic messages. The hybrids are captured  
and washed at high stringency using streptavidin coupled  
to paramagnetic particles and a magnetic separation stand.  
mRNA was selected by a single purification in this system  
25 and the yields ranged from 1-2% of the total RNA yield.  
The integrity of both the total and mRNA was analyzed by  
gel electrophoresis and ethidium bromide staining.

## cDNA Synthesis and Library Construction

cDNA was synthesized using the Superscript™ lambda  
30 system (catalog #18256-016) in conjunction with either the  
IZiplox™ vector (catalog #19643-014) or the pSV-SPORT-1  
vector (catalog #15388-010) from GibcoBrl with the

-33-

following modifications from the standard protocol. cDNA was labeled only in the first or second strand (but not both) with  $\alpha^{32}\text{p}$  dCTP and crude estimates of quantity were made by inspection of ethidium bromide staining of  
5 aliquots of cDNA fractions.

Macaque Expression Library: The size fractionation procedure was also modified slightly for construction of the macaque expression library to ensure large (>1.5kb) inserts. After one round of fractionation, only the first  
10 (largest) fraction of cDNA was saved and the remaining fractions were pooled and subjected to a subsequent round of fractionation. The top fraction from the next round was pooled with the top fraction from the previous round and the second fraction from this round was also used.  
15 These two fractions were precipitated and put into ligations with the pSV-SPORT-1 vector, and a fraction of each ligation was transformed into electrocompetent DH10B bacteria (Gibco) to estimate both the titer of the library and the average insert size. Estimates from ligation of  
20 only the top largest cDNA fraction revealed the potential of making up 2.4 million independent clones with an average insert size of 1.9kb and a median size of 2kb.

The actual library screened consisted of 150,000 independent clones which were plated at a density of 1,500  
25 clones/plate on 100 LB agar plates (to generate 100 pools of 1,500 clones/pool) with ampicillin at 50  $\mu\text{g/ml}$  and grown overnight at 37°C. For purification of individual pools, each plate was overlaid with approximately 2 ml of luria broth (LB), the colonies were scraped off each plate  
30 with a standard tissue culture cell scraper, and bacterial suspensions were transferred to microfuge tubes. Prior to purification, a glycerol stock was generated from each

-34-

pool. Plasmid DNAs were purified by use of QIAprep spin columns (QIAGEN) according to manufacturer's instructions.

#### Transfections and Adhesion Assays

CHO/P cells were seeded into 24-well plates approximately 24 hours prior to transfection at a density of 40,000 cells/well. DNAs were transiently transfected using the LipofectAMINE™ reagent (GIBCO catalog #18324-012), essentially following the recommended protocol with further optimization for 24-well plates as follows: 200 ng of DNA (representing either a plasmid pool or purified control DNAs) was diluted to 20  $\mu$ l with Opti-MEM 1 reduced serum media (GIBCO) and diluted into 20  $\mu$ l of a mixture that consists of 18  $\mu$ l Opti-MEM 1 and 2  $\mu$ l of lipofectamine reagent. This liposome mixture was then incubated for approximately 30 minutes at ambient temperature, after which 200  $\mu$ l of Opti-MEM 1 was added, and the entire mixture was then overlayed onto a well of CHO/P cells and returned to the incubator. After a 2.5 hour incubation, 240  $\mu$ l of MEM- $\alpha$  (Gibco) media with 20% fetal calf serum (FCS) was added to each well, and the cells were incubated for an additional 18-24 hours. The media was then changed to standard MEM- $\alpha$  with 10% FCS, and the adhesion assay was performed approximately 20-24 hours later.

For the adhesion assays in the expression cloning screen, the murine T cell lymphoma TK1 was used to detect cDNAs mediating cell adhesion. TK1 cells were resuspended at a density of  $2 \times 10^6$ /ml in an assay buffer which consisted of HBSS (without  $\text{Ca}^{++}$  and  $\text{Mg}^{++}$ ) supplemented with 2% bovine calf serum, 20 mM HEPES pH 7.3, 2 mM  $\text{Mg}^{++}$  and 2 mM  $\text{Ca}^{++}$ . Each well transfected with a DNA pool was pre-incubated with .25 ml of a combined supernatant

-35-

containing monoclonal antibodies to both human VCAM-1 (2G7; Graber et al., *J. Immunol.* 145:819 (1990)) and murine MAdCAM-1 (MECA-367; Streeter et al., *Nature* 331:41 (1988)) in order to eliminate adhesion mediated by VCAM-1 (which is expressed at high levels in primate lymph nodes) or any potential contaminating murine MAdCAM-1 expression plasmids. After incubation at 4°C for 15 minutes, .25 ml of the TK1 cell suspension ( $5 \times 10^5$  TK1 cells) was added to each well and incubation on a rocking platform was continued for an additional 30 minutes at 4°C. Plates were washed by gently inverting in a large beaker of phosphate buffered saline (PBS) followed by inversion in a beaker of PBS with 1.5% gluteraldehyde for fixation for a minimum of 1 hour. Wells were then examined microscopically (10X objective) for rosetting of TK1 cells mediated by the pools of cDNA clones.

The following plasmids were utilized for functional adhesion assays: pSV-SPORT-1 (GIBCO) or pcDNA3 (Invitrogen) controls, murine MAdCAM-1 in pCDM8 (pCDMAD-7), seven domain human VCAM-1 in pcDNA3 (pCD3VCAM), and human MAdCAM-1 in pcDNA3 (pCDhuMAd4). Monoclonal antibodies used were anti-murine MAdCAM-1 MECA-367, anti-human VCAM-1 2G7, anti-murine  $\alpha 467$  DATK32, anti-murine  $\beta 7$  FIB 504, anti-human  $\alpha 467$  ACT-1, anti-human integrin  $\beta 1$  (CD29) from Becton Dickinson (catalog #550034) and murine IgG1 and rat IgG2A as irrelevant controls (Sigma, St. Louis, MO).

Functional adhesion assays used the following cell lines: the murine T cell lymphoma TK1, RPMI 8866, a human B cell lymphoma which expresses  $\alpha 467$  (and not  $\alpha 461$ ) and JURKAT, a human T cell line which expresses  $\alpha 461$  (and not  $\alpha 467$ ). Assays were performed by transient transfection of plasmids encoding LERK-2 (also known as Ephrin B2),

-36-

various species of MADCAM-1, human VCAM-1, and control plasmids into CHO/P cells as described above with the following exception: as several wells were to be transfected for antibody inhibition studies, a master liposome mix with multiples of the wells to be transfected was first made for each plasmid. This ensured that the same liposome mixture was transfected into each well. On the day of the assay, monoclonal antibodies were incubated with cells at 20  $\mu$ g/ml at 4°C for 15 minutes prior to the start of the assay. Supernatants of 2G7, MECA 367 and DATK 32 were all used either neat or diluted 1:2. Assays were fixed as described above and quantitated by counting both lymphocytes and CHO cells in a field at 20X magnification. For each assay, the number of lymphocytes bound per CHO/P cell was averaged as a minimum of four fields with standard error. Results in each case are from one of three experiments performed with similar results.

#### Purification of Primate Clones

Pools yielding one or more TK1 rosettes were further subfractionated by the following protocol: DNA representative of a positive pool was retransformed into DH10B and plated on 96 plates at a density of approximately 200 colonies/plate. Nitrocellulose filters were used to generate replica plates, and one set of each plate was then subjected to DNA purification and subsequent adhesion assays as described above. A replica plate representative of a positive pool was then further subfractionated into pools of 5 colonies which were replica plated and grown overnight in LB AMP media. After one more round of DNA purification and adhesion assays, individual clones were then grown up and the clones conferring adhesion of the TK1 cells were identified.

-37-

## DNA Sequencing

Plasmids used in sequencing were as follows: The entire macaque LERK-2 cDNA was first isolated in the original vectors pSV-SPORT-1 and pZL1 (rescued from IZiplox™), respectively. Based on restriction mapping, subcloned fragments into Stratagene's Bluescript® vectors were also made to facilitate sequencing from internal regions of the cDNAs. After initial sequence analysis of these clones, oligonucleotide primers were made to complete the sequence. Overlapping sequence of both strands was obtained to ensure fidelity of the reported sequence. Sequence analysis utilized the SEQUENASE™ 7-deaza-dGTP DNA sequencing kit with SEQUENASE version 2.0 T7 DNA polymerase (United States Biochemical) and 35SdCTP (Amersham Life Science and New England Nuclear). Sequences were entered and analyzed using the Lasergene system (DNASTAR Inc.).

Sequence alignments were performed by the Clustal method (part of the Lasergene program) using a gap penalty of 10 and a gap length penalty of 10. Pairwise alignment parameters were: ktuple=2, gap penalty=5, window=4 and diagonals saved=4

## Northern Blot Analysis

Northern blots used were human multiple tissue northern I and II (commercially prepared by Clontech). Blots were pre-hybridized at 68°C for 1 hour in ExpressHyb (Clontech). cDNA's were labeled with  $\alpha^{32}\text{P}$ -dCTP by priming with random hexamers. Hybridization was performed at 68°C for 1 hour in ExpressHyb with denatured probe at a concentration of  $2 \times 10^6$  CPM/ml. Blots were then washed for 20 minutes in 2XSSC, .05% SDS at room temperature, followed by high stringency washes at 50°C, 60°C, or 65°C



-38-

in .1X SSC, .1% SDS for 20 minutes per wash and exposed to Kodak XAR film with an intensifying screen.

#### Stable LERK-2 Transfectants

The mouse L1-2 cell line is derived from a pre-B lymphoma, and was obtained from Dr. Eugene Butcher (Stanford University, Stanford, CA). The gene encoding the macaque cDNA for LERK-2 was subcloned into the pcDNA3 vector (Invitrogen), linearized by digestion with the restriction enzyme ScaI (Gibco), and transfected by the following protocol: L1-2 cells were grown to a density of approximately  $10^6$ /ml. Either 50, 25 or 12.5 million cells were washed in HBSS and then resuspended in .8 ml of a buffer consisting of Hanks balanced salt solution supplemented with 20 mM HEPES, pH 7.05. A solution consisting of 20  $\mu$ g of linearized plasmid, 500  $\mu$ g of tRNA and HBSS to bring the final volume to 200  $\mu$ l was added to the cell suspension to bring the total volume to 1 ml. After a 10-minute incubation at room temperature, the cell/DNA mixture was transferred to an electroporation cuvette (BioRad, Richmond, CA) and electroporated at 250 volts, 960 mF in a BioRad gene pulser. Following another 10-minute incubation, the cells were diluted to 25 ml in standard L1-2 growth media (RMPI 1640, 10% Hyclone fetal bovine serum, Penicillin/Stytreptomycin (50 U/ml [Gibco]) and L Glutamine (0.29 mg/ml [Gibco])) and returned to the incubator. Forty-eight hours later, the cells were pelleted by centrifugation and resuspended in 50 ml of L1-2 media supplemented with G418 (Genticin [Gibco]) at 0.8 mg/ml. Dilutions of the cell suspension were plated in 96-well microtiter plates and single colonies were grown up analyzed for expression of LERK-2.

-39-

Clones expressing LERK-2 could be detected by aggregation with TK1 cells. L1-2 (non-transfected cells) and TK1 cells both grow as single cell suspensions. Surface expression of LERK-2 is inferred by its ability to mediate aggregation by virtue of its interaction with the putative EPH receptor in on TK1 cells (which we now know is Nuk/EphB2). RNA was prepared from L1-2 lines that mediated significant aggregation, and RNA PCR confirmed that these lines did in fact contain message for LERK-2; as a control, L1-2 cells which did not aggregate were negative by this PCR analysis.

CHO cells stably transfected with LERK-2 cDNA were prepared as described above with the following exceptions. Media for CHO cell growth was  $\alpha$ -MEM with deoxyribonucleosides (Gibco) and 10% fetal calf serum (Gibco) and Penicillin/Streptomycin (50 U/ml [Gibco]) and L Glutamine (0.29 mg/ml [Gibco]). Selection media consisted of the same media with 0.55 mg/ml G418 (Gibco). Single clones were grown up and analyzed for their ability to mediate adhesion of TK1 cells. Using this criteria, a line called CHO5C2 was established.

#### Monoclonal Antibodies

Monoclonal antibodies against primate LERK-2 were generated by immunizing C57BL/6 mice with L1-2 LERK-2 transfectants. Mice were immunized intraperitoneally with 10 million cells, resuspended in HBSS 3 times at two-week intervals, and a final immunization was injected intravenously.

A successful fusion was performed which generated approximately 5,000 hybridomas. Four days after the final intravenous injection, the spleen was removed and a single cell suspension was prepared in serum-free DMEM media.

-40-

These cells were fused with the fusion partner SP2/0, according to the method of Galfre et al. (Nature 299:550-552 (1977)). 20 ml of spleen cells and 20 ml of SP2/0 cells were combined, spun at 800g for 5 minutes, and the media was aspirated. A solution of 50% polyethylene glycol 1500 (PEG 1500) (Boehringer Mannheim, Indianapolis, In) prewarmed to 37°C was added to the cell pellet over 2 minutes, followed by 10 ml of DMEM media over 3 minutes. The cell suspension was spun at 600g for 3 minutes and the supernatant removed. The pellet was resuspended gently in DMEM media containing 20% fetal calf serum, 2 mM L-glutamine, 100 U/ml penicillin, 100 µg/ml streptomycin sulfate, and HAT selection media (Sigma). Cells were plated into 10 96-well flat bottom microtiter plates at 200 µl/well.

Ten days after the fusion, supernatants from the wells were screened for reactivity against CHO LERK-2 transfectants with the CHO5C2 line as a positive control using a fluorescently-labeled (FITC) anti-mouse antibody (Jackson Labs). Several antibodies were selected for strong reactivity against the CHO LERK-2 transfectants. The putative positive antibodies were then screened in a plus minus screen using non transfected CHO cells as a negative control. Two antibodies, designated 4A1 and 2A1, were confirmed as anti LERK-2 mAbs.

These supernatants were subsequently screened for their ability to block adhesion of TK1 cells to CHO LERK-2 transfectants, and both mAbs could inhibit adhesion of TK1 cells to LERK-2 transfectants. Inhibition experiments were performed essentially as described for supernatants above. These two blocking hybridomas were subcloned using limiting dilution.

-41-

The isotype of the two mAbs 2A1 and 4A1 was determined to be IgG2A and IgG1, respectively, by the SBA Clonotyping System/HRP (Southern Biotechnology Associates, Inc.). Maxisorb plates were first coated with a trapping  
5 antibody, goat anti-mouse Ig, at 5  $\mu$ g/ml overnight. The plates were then washed 3 times with PBS/Tween and blocked with 1% BSA for 1 hour at room temperature. After another 3 washes with PBS/Tween, antibodies to be tested and purified isotype standards were added at 100  $\mu$ g/ml and  
10 incubated for 2 hours at room temperature. Then the plates were washed and the mAbs and isotype standards were incubated with a variety of HRP-conjugated anti-isotype antibodies for 1 hour at room temperature followed by detection with OPD substrate.

15 FACS analysis

100  $\mu$ l of whole blood or cell suspensions were added to tubes containing specific (4A1) or irrelevant (IgG1, MOPC21, Sigma, St. Louis, MO) primary antibodies in blocking solution to a final concentration of 10  $\mu$ g/ml.  
20 Cell suspensions were incubated at 4°C for 20 minutes, washed twice with PBS and subsequently incubated with secondary antibody (goat anti mouse-PE, Jackson ImmunoResearch, West Grove, PA) at a final concentration of 10 mg/ml in blocking solution at 4°C for 15 minutes.  
25 Subsequently, 2 ml of FACS-lysis solution (Becton-Dickinson) was added, and tubes were incubated for 10 minutes at room temperature. Following two final PBS washes, cells were resuspended in 300 ml PBS with 1% fetal calf serum and stored at 4°C until analyzed.  
30 FACS analysis: Two-color analysis was performed on a Becton Dickinson FACScan flow cytometer. Gating for lymphocytes and granulocytes was based upon forward and

-42-

side scatter characteristics. For each sample, a minimum of 10,000 cells was counted.

## RESULTS

### FACS analysis

5       The two anti-LERK-2 mAbs were examined for cross reactivity to human LERK-2 by FACS analysis of human umbilical vein endothelial cells (HUVECs) in the presence or absence of TNF- $\alpha$ . While the 2A1 mAb failed to react, 4A1 did react and was slightly induced by a both 4 and 24  
10   hour incubation with TNF- $\alpha$ . This is consistent with the results of Beckmann et al. (*EMBO J.* 13:3757-3762 (1994)) which showed expression of LERK-2 RNA in northern blots made from TNF- $\alpha$ - stimulated HUVECS

      The mAb 4A1 was also used to investigate the  
15   expression of LERK-2 on leukocytes by FACS analysis. Peripheral blood mononuclear cells (PBMCs) were isolated from human blood using Ficoll. LERK-2 expression on subsets of PBMCs were examined by 2-color staining using both 4A1 and another marker for a subset of leukocytes.  
20   It was discovered that LERK-2 is highly expressed on monocytes/macrophages (CD14 positive cells). Some levels of expression were also seen on lymphocytes. The T cell subsets that are positive for LERK-2 include CD4+, CD8+, and CD45RO+ (Figures 1A-C).

25       To further confirm the expression of LERK-2 on leukocyte subsets, a variety of monocytic and lymphocytic cell lines were studied by FACS using 4A1 as a staining reagent. The cell lines include HL60, HSB-2, HUT78, IBW4, JY, KCA K562, RAMOS, Raji, RPMI8866, and THP-1 cells.  
30   Among these cell lines, THP-1 and HSB-2 express a high level of LERK-2. HL60, HUT78, KCA, K562, RAMOS, and Raji express moderate levels of LERK-2, while IBW4, JY and RPMI

-43-

do not express LERK-2 by this FACS analysis (Figures 2A-B). This result, showing the high level expression of LERK-2 on a monocytic cell line (THP-1) and a lymphocytic cell line (HSB-2), is consistent with the expression  
5 pattern on PBMC.

The expression of LERK-2 on PBMCs and leukocyte cell lines was confirmed by RT-PCR (Figure 3). Whole cell polyadenylated RNA was isolated from PBMC and the leukocyte cell lines used in the FACS analysis using  
10 Oligotex Direct mRNA Isolation Kit (QIAGEN). Cells were lysed and the lysates were incubated with oligo(dT)<sub>n</sub>-coated latex beads to select for polyA<sup>+</sup> RNA. The Oligotex beads were wash and polyA<sup>+</sup> RNA was eluted from the beads. Reverse transcription (RT) was performed in the presence  
15 of MuLV reverse transcriptase, dNTP, Rnase inhibitor, Oligo(dT)<sub>16</sub>, and appropriate polyA<sup>+</sup> RNA as template. The reaction was carried out at 42°C for 1 hour followed by heat inactivation of the enzyme. Polymerase chain reaction (PCR) was performed using a pair of primers  
20 specific for LERK-2, the forward amplification primer 5'-GCCCCGAGCAGAAGCA-3' (SEQ ID NO: 1) and the reverse amplification primer  
5'-CAGGAAGATGATGATGAGCAG-3' (SEQ ID NO: 2), in the presence of the RT product. PCR product specific for  
25 LERK-2 (600 bp) was obvious in all lines that were brightly positive for LERK-2 including PBMC, THP-1, and HSB-2 cells.

#### Construction of LERK-2-Ig chimera and FACS analysis

A chimera containing the ectodomain of LERK-2 and the  
30 Fc domain of human IgG1 was constructed as described previously (Tidswell et al., *J. Immunol* 159:1497-1505 (1997)) using the forward amplification primer 5'-

-44-

GGAGCTTCCACCATGGCTCGGCCTGGGCAG-3' (SEQ ID NO: 3) and the reverse amplification primer  
5'-GGACTAGTGCCACCTTTGAGTTGAAGAAG-3' (SEQ ID NO: 4) to isolate the LERK-2 ECD. The construct was transiently  
5 transfected into CHO-P cells using Lipofectamine (GIBCO), and the supernatant of the transfected cells was harvested. This supernatant was used as a FACS reagent to stain leukocyte and endothelial cell lines. The leukocyte cell lines tested include TK1, HL60, HSB2, Raji,  
10 RAMOS, RPMI, JY, and THP-1. Among the leukocyte cell lines, only TK1 cells are positive when stained with LERK-2-Ig chimera, indicating that TK1 cells express a receptor for LERK-2, which is consistent with other results described herein (Figure 4).

15 The expression of receptor for LERK-2 on HUVEC cells was also investigated in the presence or absence of TNF- $\alpha$  stimulation. The data showed that a receptor for LERK-2 is expressed on HUVEC cells, while TNF- $\alpha$  stimulation does not increase the expression of this receptor on HUVECs  
20 (Figure 5).

#### Summary of LERK II Tissue Staining

The expression of LERK-2 in normal and inflamed human tissues by immunohistochemistry and flow cytometry (see Table). Human tissues (normal and inflamed) were obtained  
25 from the National Disease Research Institute, a service organization funded by the National Institutes of Health.

Immunohistochemical analysis for LERK-2 was performed on frozen tissue samples. Briefly, tissue was sectioned at a thickness of 4 mm, desiccated, and then fixed in 2%  
30 paraformaldehyde/0.5 X PBS for 10 minutes at 4°C. After PBS washing, nonspecific antibody binding sites were blocked with 10% normal goat serum/5% human AB serum/PBS

-45-

for 30 minutes at room temperature. Next the purified, LERK-2 murine antibody (4A1) was diluted to a concentration of 10 mg/ml in 0.3% Triton X 100/0.2% Tween 20/1% FCS/5% human AB serum/0.1% sodium azide and applied to tissue sections overnight at 4°C. An isotype-matched irrelevant monoclonal antibody (IgG1, MOPC21) was used as a negative control on step sections of tissues. Subsequently, biotinylated goat anti-mouse IgG and avidin-biotin-alkaline phosphatase complexes (Biogenex, San Ramon, CA) were added in sequence. Fast Red (Biogenex, San Ramon, CA), containing 2% levamisole to block endogenous alkaline phosphatase activity, was used as the chromogen and Mayers hematoxylin as the counterstain.

LERK-2 is widely expressed on vascular endothelium in all organs examined. The staining intensity is quite variable between tissues. In tissues, patchy staining of vascular smooth muscle is frequently observed. Sections of lymph node reveal staining on mononuclear cells (lymphocytes, monocyte/macrophages) within the subcapsular and medullary sinuses as well as the paracortex. In sections of inflamed colon, LERK-2 staining is sometimes more intense on endothelia of vessels adjacent to regions of inflammation.

While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the spirit and scope of the invention as defined by the following claims.



-46-

## CLAIMS

What is claimed is:

1. A method of modulating LERK-2-mediated cell adhesion,  
comprising the step of contacting a cell selected  
5 from the group consisting of:
  - a) a cell expressing LERK-2;
  - b) a cell expressing a receptor for LERK-2with an inhibitor or promoter of LERK-2-mediated cell adhesion.
- 10 2. A method according to Claim 1, wherein the receptor for LERK-2 is selected from the group consisting of Nuk, Cek 5, Tyro 5 and ERK.
3. A method according to Claim 2, wherein the receptor for LERK-2 is Nuk.
- 15 4. A method according to Claim 1, wherein the cell expressing LERK-2 is a cell which bears LERK-2 in nature.
5. A method according to Claim 1, wherein the cell  
expressing a receptor for LERK-2 is a cell which  
20 bears a receptor for LERK-2 in nature.
6. A method according to Claim 1, wherein the cell expressing a receptor for LERK-2 is a lymphoid cell.
7. A method according to Claim 6, wherein the cell expressing a receptor for LERK-2 is a T cell.

-47-

8. A method of inhibiting LERK-2-mediated cell adhesion, comprising the step of contacting a cell selected from the group consisting of:
- a) a cell expressing LERK-2; and
  - 5 b) a cell expressing a receptor for LERK-2
- with an inhibitor of LERK-2-mediated cell adhesion selected from the group consisting of:
- a) an antibody or functional antibody fragment which inhibits LERK-2-mediated cell adhesion;
  - 10 b) soluble LERK-2; and
  - c) LERK-2-Ig chimera.
9. A method according to Claim 8, wherein the inhibitor is an antibody or functional antibody fragment which inhibits LERK-2-mediated cell adhesion.
- 15 10. A method according to Claim 9, wherein the antibody or functional antibody fragment is an anti-LERK-2 antibody or fragment selected from the group consisting of 2A1, a functional antibody fragment of 2A1, 4A1 and a functional antibody fragment of 4A1.
- 20 11. A method according to Claim 1, wherein the promoter is selected from the group consisting of:
- a) isolated LERK-2;
  - b) LERK-2 chimeras; and
  - c) LERK-2 mimics.
- 25 12. A method of detecting or identifying an inhibitor of LERK-2-mediated cell adhesion, comprising the steps of:
- a) combining an agent to be tested with a composition comprising a cell expressing LERK-2

-48-

and a composition comprising a cell expressing a receptor for LERK-2 under conditions suitable for binding of LERK-2 to a receptor for LERK-2; and

- 5        b) detecting or measuring cell adhesion between said cell expressing LERK-2 and said cell expressing a receptor for LERK-2, whereby inhibition of cell adhesion by the agent is indicative that the agent is an inhibitor.
- 10    13. A method according to Claim 12, wherein the receptor for LERK-2 is selected from the group consisting of Nuk, Cek 5, Tyro 5 and ERK.
14. A method according to Claim 13, wherein the receptor is Nuk.
- 15    15. A method according to Claim 12, wherein cell adhesion between a cell expressing LERK-2 and a cell expressing a receptor for LERK-2 can mediate cellular signalling and/or a cellular response, and cell adhesion is monitored by detecting or measuring a
- 20       signalling activity or cellular response of one or both of the cells in response thereto.
16. An inhibitor of LERK-2-mediated cell adhesion identified according to the method of Claim 12.
17. An inhibitor according to Claim 16, wherein the
- 25       inhibitor is an antibody or functional antibody fragment.

-49-

18. A method of detecting or identifying a promoter of LERK-2-mediated cell adhesion comprising the steps of:
- 5 a) combining an agent to be tested with a composition comprising a cell expressing LERK-2 and a composition comprising a cell expressing a receptor for LERK-2 under conditions suitable for binding of LERK-2 to a receptor for LERK-2; and
- 10 b) detecting or measuring cell adhesion between said cell expressing LERK-2 and said cell expressing a receptor for LERK-2, whereby induction or stimulation of cell adhesion by the agent is indicative that the agent is a
- 15 promoter.
19. A promoter of LERK-2-mediated cell adhesion identified according to the method of Claim 18.
20. A method according to Claim 18, wherein cell adhesion between a cell expressing LERK-2 and a cell
- 20 expressing a receptor for LERK-2 can mediate cellular signalling and/or a cellular response, and cell adhesion is monitored by detecting or measuring a signalling activity or cellular response of one or both of the cells in response thereto.
- 25 21. An antibody or functional antibody fragment which inhibits LERK-2-mediated cell adhesion.
22. The antibody or functional antibody fragment of Claim 21, wherein the antibody or fragment inhibits binding of LERK-2 to a receptor for LERK-2.

-50-

23. The antibody or functional antibody fragment of Claim 22, wherein said antibody or antibody fragment is an anti-LERK-2 antibody.
- 5 24. An antibody or functional antibody fragment according to Claim 23, wherein the antibody is selected from the group consisting of monoclonal antibody 2A1, monoclonal antibody 4A1, a functional antibody fragment of 2A1 and a functional antibody fragment of 4A1.
- 10 25. An antibody or functional antibody fragment according to Claim 21, wherein said antibody or fragment can compete with monoclonal antibody 2A1 or monoclonal antibody 4A1 for binding to LERK-2 or portion thereof.
- 15 26. An antibody or functional antibody fragment according to Claim 21, wherein said antibody or antibody fragment selectively inhibits the interaction of LERK-2 with Nuk.
- 20 27. An antigen binding fragment of an antibody according to Claim 21.
28. Monoclonal antibody 2A1 or an antibody having an epitopic specificity which is the same as or similar to that of 2A1.
- 25 29. Monoclonal antibody 4A1 or an antibody having an epitopic specificity which is the same as or similar to that of 4A1.

-51-

30. The hybridoma cell line deposited under ATCC  
Accession No. HB-12412.
31. The hybridoma cell line deposited under ATCC  
Accession No. HB-12413.
- 5 32. A monoclonal antibody or antigen binding fragment  
thereof produced by the hybridoma cell line of Claim  
30.
33. A monoclonal antibody or antigen binding fragment  
thereof produced by the hybridoma cell line of Claim  
10 31.
34. A method for treating an angiogenic disease,  
comprising administering to a mammal a  
therapeutically effective amount of an inhibitor of  
LERK-2-mediated cell adhesion, whereby angiogenesis  
15 is reduced.
35. A method according to Claim 34, wherein the inhibitor  
is an antibody or functional antibody fragment which  
inhibits binding of LERK-2 to a receptor for LERK-2.
36. A method according to Claim 35, wherein said antibody  
20 or antibody fragment is an anti-LERK-2 antibody.
37. A method according to Claim 36, wherein the antibody  
or antibody fragment is selected from the group  
consisting of monoclonal antibody 2A1, monoclonal  
antibody 4A1, a functional antibody fragment of 2A1  
25 and a functional antibody fragment of 4A1.

-52-

38. A method according to Claim 36, wherein said antibody or fragment can compete with monoclonal antibody 2A1 or monoclonal antibody 4A1 for binding to LERK-2 or portion thereof.
- 5 39. A method according to Claim 35, wherein said antibody or antibody fragment inhibits the interaction of LERK-2 with Nuk.
40. A method for treating an inflammatory disease or condition, comprising administering to a mammal a  
10 therapeutically effective amount of an inhibitor of LERK-2-mediated cell adhesion, whereby inflammation is reduced.
41. A method according to Claim 40, wherein the  
15 inflammatory disease or condition is a T cell-mediated disease or condition.
42. A method according to Claim 40, wherein the inhibitor is an antibody or functional antibody fragment which inhibits binding of LERK-2 to a receptor for LERK-2.
43. A method according to Claim 42, wherein said antibody  
20 or antibody fragment is an anti-LERK-2 antibody.
44. A method according to Claim 43, wherein the antibody or antibody fragment is selected from the group consisting of monoclonal antibody 2A1, monoclonal antibody 4A1, a functional antibody fragment of 2A1  
25 and a functional antibody fragment of 4A1.

-53-


45. A method according to Claim 43, wherein said antibody or fragment can compete with monoclonal antibody 2A1 or monoclonal antibody 4A1 for binding to LERK-2 or portion thereof.
- 5 46. A method according to Claim 42, wherein said antibody or antibody fragment inhibits the interaction of LERK-2 with Nuk.
47. A method of antitumor therapy, comprising administering to a mammal a therapeutically effective  
10 amount of an inhibitor of LERK-2-mediated cell adhesion, whereby angiogenesis is reduced, thereby inhibiting tumor growth.
48. A method for enhancing angiogenesis, comprising  
15 administering to a mammal a therapeutically effective amount of a promoter of LERK-2-mediated cell adhesion, whereby angiogenesis is enhanced.
49. A method according to Claim 48, wherein the promoter  
20 is selected from the group consisting of isolated LERK-2, LERK-2 chimeras, and LERK-2 mimics.



1/8

## Immunostaining with 4A1 and anti-CD4

Gate on lymphocytes



Gate on monocytes








FIG. 1A

SUBSTITUTE SHEET (RULE 26)

IMMUNOSTAINING WITH 4A1 AND ANTI-CD14




FIG. 1B




FIG. 1C

## Immunostaining of cell lines




FIG. 2A

SUBSTITUTE SHEET (RULE 26)

5/8










**K562****RAMOS****Raji****RPMI 8866****THP-1****HUVEC****HUVEC+TNF****FIG. 2B****SUBSTITUTE SHEET (RULE 26)**

FIG. 3




Best Available Copy


7/8




TK1




HL60




HSB2




Raji




RAMOS



RPMI



JY



THP1

FIG. 4

8/8





FIG. 5



## SEQUENCE LISTING

<110> LeukoSite, Inc.  
Briskin, Michael J.  
Zou, Lily

<120> MODULATION OF LERK-2-MEDIATED CELL  
ADHESION

<130> LKS96-06pA

<150> US 60/060,757

<151> 1997-10-02

<160> 4

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer Sequence

<400> 1

gcccccgagc agaagca

17

<210> 2

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer Sequence

<400> 2

caggaagatg atgatgagca g

21

<210> 3

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Primer Sequence

<400> 3

ggagcttcca ccatggctcg gcctgggcag

30

2/2

<210> 4  
<211> 29  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Primer Sequence

<400> 4  
ggactagtgc cacctttgag ttgaagaag

29

SUBSTITUTE SHEET (RULE 26)